

Selected Publications by Edward Wise, Ph.D.

Wise, E.A., (in press). Selected MMPI-2 Scores of Forensic Offenders in a Community Setting. *Assessment*.

Wise, E.A. (2005). Effectiveness of intensive outpatient programming in private practice: Integrating practice, outcomes and business. *American Psychologist*, 60(8), 885-895.

Wise, E. A. (2004). Methods for analyzing psychotherapy outcomes: A review of clinical significance, reliable change and recommendations for future directions. *Journal of Personality Assessment*, 82(1), 50-59.

Wise, E. A. (2003). Personality style codetype concordance between the MCMI and MBHI. *Journal of Clinical Psychology* (Special Issue: Coping, Part 2-Measurement), 59(12), 1279-1294. (Re-published: original work published 1994).

Wise, E. A. (2003). Psychotherapy outcome and satisfaction methods applied to Intensive Outpatient Programming in a private practice setting. *Psychotherapy: Theory, Research and Practice*, 40(3), 203-214.

Wise, E.A. (2003). Empirical validation of a mental health Intensive Outpatient Program in a private practice setting. *American Journal of Orthopsychiatry*, 73(4), 405-410.

Wise, E.A. (2003). A private practice model for intensive outpatient services. *Psychiatric Services*, 54(4), 572 - 573.

Wise, E.A. (2002). Relationships of personality disorders with MMPI-2 malingering, defensiveness and inconsistent response scales among forensic examinees. *Psychological Reports*, 90, 760 - 766.

Wise, E.A. (2001). PPO primary care physician survey on mental health utilization. *Journal of Psychotherapy in Independent Practice*, 2(4), 57-72.

Wise, E.A. (2001). The comparative validity of MMPI-2 and MCMI-II personality disorder classifications with forensic examinees. *Journal of Personality Disorders*, 15(3), 275-279.

Wise, E.A. (2000). Mental health intensive outpatient programming: An outcome and satisfaction evaluation of a private practice model. *Professional Psychology: Research and Practice*, 31(4), 412-417.

Wise, E.A. (1999). Clinical outcomes, client satisfaction and innovative programming in private practice. In Coughlin, K.M. (Ed). *Behavioral Outcomes & Guidelines Sourcebook: A Practical Guide to Measuring, Managing and Standardizing Mental Health and Substance Abuse Treatment* (pps. 135-143). Faulkner & Gray, NY, NY.

Wise, E. A. (1998). Group Therapy Needs Assessment with Community Employee Assistance Programs. *Professional Psychology: Research and Practice*, Vol. 29, No. 1, 56-58.

Wise, E.A. (1996). Diagnosing Post-Traumatic Stress Disorder with the MMPI clinical scales: A review of the literature. *Journal of Psychopathology and Behavioral Assessment*, 18(1), 71-82.

Wise, E.A. (1996). Comparative validity of the MMPI-2 and MCMI-II personality disorder classifications. *Journal of Personality Assessment*, 66(3), 569-582.

Wise, E.A. (1996). Guidelines for interpreting MMPI PTSD profiles. *Anxiety Disorders Practice Journal*, 2(2), 161-170.

Wise, E.A. (1995). Personality disorder correspondence between the MMPI, MBHI and MCMI. *Journal of Clinical Psychology*, 51(6), 790-798.

Wise, E.A. (1994). Managed care and the psychometric validity of the MMPI and MCMI personality disorder scales. *Psychotherapy in Private Practice*, 13(3), 81-97.

Wise, E.A. (1994). Projective identification and countertransference in group psychotherapy. *Voices: The Art and Science of Psychotherapy*, 30(2), 40-45.

Wise, E.A. (1994). Personality disorder correspondence between the MBHI and the MMPI. *Journal of Clinical Psychology in Medical Settings*, 1(2), 167-181.

